
Stripped Star plus Compact Object Binaries: Identifying the Progenitors of Neutron Star Mergers

Maria R. Drout University of Toronto Carnegie Observatories

Masses in the Stellar Graveyard


LIGO-Virgo-KAGRA Black Holes LIGO-Virgo-KAGRA Neutron Stars



LIGO-Virgo-KAGRA | Aaron Geller | Northwestern


The Evolutionary Path to a BNS

There are multiple stages with (large) uncertainties




What are Stripped Stars? How Can They Help?

What are Stripped Stars? Hydrogen-rich envelopes removed via binary interaction




Common Envelope Evolution




What are Stripped Stars?

Hydrogen-rich envelopes removed via binary interaction



Stripped cores are hot and compact

The Evolutionary Path to a BNS Multiple Phases of Envelope Stripping are Required



They probe of the outcomes of the common envelope stage and the physical configuration of the system prior to the second supernovae






Image courtesy of T. Tauris

They probe of the outcomes of the common envelope stage and the physical configuration of the system prior to the second supernovae

They can be multimessenger sources themselves!



Y. Götberg et al. (2020)




Y. Götberg et al. (2020)

They probe of the outcomes of the common envelope stage and the physical configuration of the system prior to the second supernovae

They can be multimessenger sources themselves!


GW source	Number	Frequency	Chirp mass	Distance
Stripped star + NS^a	$\sim 0 - 4^* \text{ (SNR} > 5)$	< 0.2 mHz	$1.6 - 2.6 \ M_{\odot}$	≲ 1 kpc
Stripped star + WD^a	$\sim 0 - 100^* \text{ (SNR} > 5)$	< 0.6 mHz	$0.3-1.6~M_{\odot}$	$\lesssim 1 \text{ kpc}$




Y. Götberg et al. (2020)

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions



- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions



There are currently no confirmed stripped star + NS/BH binaries

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions




Image courtesy of T. Tauris

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

Local Group Galaxies & Galactic Plane

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions

Where to Look?

Local Group Galaxies & Galactic Plane

(large area, dense environments)

Image Credit: NASA/Swift/S. Immler (Goddard) and M. Siegel (Penn State)

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?


- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Stripped Star

Accretion



- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?


Stripped Star

Accretion

Ultraviolet is key

Stripped Star Compact Object Binaries:


What (specific) time-domain/multi-wavelength observations?



Y. Gotberg et al. (2016, 2018)

Stripped Star Compact Object Binaries:

What (specific) time-domain/multi-wavelength observations?



ions

Accretion


Drout, Götberg, et al. submitted

Stripped Star Compact Object Binaries: Proof of Concept Search in the Magellanic Clouds




Image Credit: NASA/Swift/S. Immler (Goddard) and M. Siegel (Penn State)

Stripped Star Compact Object Binaries: Proof of Concept Search in the Magellanic Clouds



Stripped Star Compact Object Binaries: Proof of Concept Search in the Magellanic Clouds



A subset of stars have spectra consistent with models for "isolated" helium stars. They have "dark" companions

Götberg, Drout, et al. submitted

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Stripped Star

Accretion




Wide-Field High-Resolution UV Imager

- 1. To identify a population

Where to Look?

For What? Stripped Star

Accretion




- 1. To identify a population

Where to Look?

For What? Stripped Star

Accretion



Sources should be (weak-ish) X-ray binaries

Stripped Star Compact Object Binaries:

What (specific) time-domain/multi-wavelength observations?





Image Credit: B. Ludwig (Toronto)

Stripped Star Compact Object Binaries:

What (specific) time-domain/multi-wavelength observations?



- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Stripped Star

Accretion

Critical to Maintain High-Sensitivity High-Resolution X-ray Capabilities

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Light Curves

Velocities

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Light Curves

Velocities


- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Light Curves

Velocities



Geier et al. (2013)


- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?


For What?

Light Curves

Velocities



Geier et al. (2013)



Y. Götberg et al. (2020)


- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Light Curves

Velocities



Y. Götberg et al. (2020)

- 1. To identify a population
- 2. To determine period and mass distributions
- To determine mass loss rates and surface compositions

Where to Look?

For What?

Light Curves

Velocities

(UV) High Cadence Monitoring Moderate Resolution Spectroscopic Monitoring

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions

Where to Look?

For What?

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions


Where to Look?


For What?

Spectral (Wind) Features

Stripped Star Compact Object Binaries:


What (specific) time-domain/multi-wavelength observations?





Stripped Star Compact Object Binaries:

What (specific) time-domain/multi-wavelength observations?



Götberg, Drout, et al. submitted

- 1. To identify a population
- 2. To determine period and mass distributions
- 3. To determine mass loss rates and surface compositions

Where to Look?

For What?

Spectral (Wind) Features

Moderate resolution UV spectroscopy is critical

Stripped Star Compact Object Binaries: What type of instruments/missions?

Wide-Field High-Resolution UV Imager

High-Sensitivity High-Resolution X-ray Capabilities

(UV) High Cadence Monitoring

Moderate Resolution Spectroscopic Monitoring

Moderate resolution UV spectroscopy

Stripped Star Compact Object Binaries: What type of instruments/missions?

Wide-Field High-Resolution UV Imager


High-Sensitivity High-Resolution X-ray Capabilities


(UV) High Cadence Monitoring

Moderate Resolution Spectroscopic Monitoring

Moderate resolution UV spectroscopy

Stripped Star Compact Object Binaries: CASTOR: A Wide-Field UV Telescope





Bandpasses

Field of View

Image Quality

Observing Modes

UV (150-300nm), u' (300-400 nm), g (400-550 nm)

 $0.44 \deg \times 0.56 \deg = 0.25 \operatorname{sq.} \deg$

~0.15" FWHM

wide-field imaging; grism and 2D spectroscopy; precision photometry

Stripped Star Compact Object Binaries: What type of instruments/missions?

Wide-Field High-Resolution UV Imager

High-Sensitivity High-Resolution X-ray Capabilities

(UV) High Cadence Monitoring

Moderate Resolution Spectroscopic Monitoring

Moderate resolution UV spectroscopy